

On the concentration of the chromatic number of random graphs

Alex Scott *

June 2, 2008

Abstract

Let $0 < p < 1$ be fixed. Shamir and Spencer proved in the 1980s that the chromatic number of a random graph $G \in \mathcal{G}(n, p)$ is concentrated in an interval of length $\omega(n)\sqrt{n}$. We give an improvement on this, showing that $\chi(G)$ is concentrated in an interval of length $\omega(n)\sqrt{n}/\log n$.

1 Introduction

How concentrated is the chromatic number $\chi(G)$ of a random graph $G \in \mathcal{G}(n, p)$? For constant probability $p \in (0, 1)$, Shamir and Spencer [10] proved in the 1980s that $\chi(G)$ is concentrated in an interval of length $\omega(n)\sqrt{n}$. For sparse random graphs, much stronger concentration results are known: Shamir and Spencer [10] showed that for $p < n^{-5/6-\epsilon}$, the chromatic number is concentrated on 5 consecutive integers. Łuczak [7] sharpened this to a 2-point concentration result, while Alon and Krivelevich [2] extended 2-point concentration to the larger range $p < n^{-1/2-\epsilon}$.

The aim of this note is to show that Łuczak's approach also works for random graphs with constant density, giving a slight improvement on the concentration result of Shamir and Spencer, from $\omega(n)\sqrt{n}$ to $\omega(n)\sqrt{n}/\log n$. In particular, we prove the following theorem.

*Mathematical Institute, 24-29 St Giles', Oxford, OX1 3LB, UK; email: scott@maths.ox.ac.uk

Theorem 1. *Let $0 < p < 1$ be fixed, and suppose $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$. Then there is a function $h = h(n)$ such that, for $G \in \mathcal{G}(n, p)$, with probability $1 - o(1)$,*

$$|\chi(G) - h(n)| < \omega(n)\sqrt{n}/\log n. \quad (1)$$

Let us note that bounding the concentration of the chromatic number does not determine its likely value. The asymptotic behaviour of $\mathbb{E}\chi(G)$, where $G \in \mathcal{G}(n, p)$ and p is fixed, was resolved by Bollobás [4], while Łuczak [8] extended this result to a much wider range of values $p = p(n)$ (see also McDiarmid [9]). In the sparse case ($p = c/n$), Achlioptas and Naor [1] have given an explicit pair of consecutive integers $k, k + 1$ such that $\chi(G) \in \{k, k + 1\}$ almost surely, and Coja-Oghlan, Panagiotou, Steger [6] recently proved an explicit three-point concentration result for $p < n^{-3/4-\epsilon}$.

Finally, let us note that, for p fixed, nothing is known from below concerning concentration: it has not even been shown that the chromatic number cannot be concentrated in an interval of constant length! See Bollobás ([5] and [3]) for further discussion.

2 Proof

The proof will proceed as follows. As in Łuczak [8], we first define $h(n)$ so that $\mathbb{P}(\chi(G) \leq h(n))$ tends to 0 slowly. Then $\chi(G) > h(n)$ with probability $1 - o(1)$, so we need only bound $\chi(G)$ from above. A martingale argument shows that we can colour all but (a little more than) $\sqrt{n}$ vertices with $h(n)$ colours, so we try to colour the remaining vertices without using too many new colours. Łuczak's argument used the local sparsity of G ; here, in the dense case, we can use a fairly crude greedy algorithm, which shows that any reasonably large set S of vertices can be coloured with $O(|S|/\log n)$ colours. This will be enough to prove (1).

We first need a simple lemma on independent sets in random graphs. It is slightly cleaner to phrase it in terms of complete subgraphs.

Lemma 2. *Let $0 < p < 1$ be fixed, and suppose $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$. There is a constant $c = c(p)$ such that, for $G \in \mathcal{G}(n, p)$, with probability $1 - o(1)$, every subset $W \subset V(G)$ with $|W| > n^{1/3}$ contains a complete subgraph with at least $c \log n$ vertices.*

Proof of Lemma 2. Let $G \in \mathcal{G}(n, p)$, and suppose that $U \subset V(G)$ has $u \geq n^{1/4}$ vertices. We use Chernoff's inequality in the form that $X \sim B(r, p)$

implies $\mathbb{P}(X < rp - t) \leq \exp(-t^2/2rp)$. Thus (with $t = pu^2/5$),

$$\mathbb{P}(e(U) < pu^2/4) \leq \exp\left(-(pu^2/5)^2/2p\binom{u}{2}\right) < \exp(-pu^2/30).$$

So the probability that there is a subset of size u with fewer than $pu^2/4$ edges is at most

$$\binom{n}{u} e^{-pu^2/30} \leq \left(\frac{en}{u}\right)^u e^{-pu^2/30} = \left(\frac{en}{u} e^{-pu/30}\right)^u = o(1).$$

It follows that, with probability $1 - o(1)$, every subset of u vertices has edge density at least $p/2$; thus every subset of $u' \geq u$ vertices has edge density at least $p/2$ and hence induces a subgraph with maximal degree at least $p(u' - 1)/2$.

Suppose that this property holds. Given $W \subset V(G)$, we choose a complete subgraph greedily: set $W_0 = W$ and, for $i \geq 1$, pick $w_i \in W_{i-1}$ with $|\Gamma(w_i) \cap W_{i-1}|$ maximal and set $W_i = \Gamma(w_i) \cap W_{i-1}$, halting with the complete subgraph $\{w_1, \dots, w_i\}$ as soon as W_i is empty. The observations above imply that (for n sufficiently large) $|W_i| \geq \frac{p}{2}(|W_{i-1}| - 1) > p|W_{i-1}|/3$ whenever $|W_{i-1}| \geq u$, and so $|W_i|$ is nonempty provided $(p/2)^i < n^{-1/12}$. The result follows immediately with $c(p) = -1/12 \log(p/2)$. $\square$

Proof of Theorem 1. We may assume that $\omega(n) = o(\log \log n)$. Let $G \in \mathcal{G}(n, p)$ and let

$$h(n) = \min \left\{ r : \mathbb{P}(\chi(G) \leq r) > \frac{1}{\omega(n)} \right\}. \quad (2)$$

We shall show that (1) holds with this h .

Clearly $\mathbb{P}(\chi(G) < h(n)) \rightarrow 0$ as $n \rightarrow \infty$; thus it suffices to show that, with probability $1 - o(1)$, we have $\chi(G) < h(n) + \omega(n)\sqrt{n}/\log n$. Let

$$s(G) = \max\{|W| : W \subset V(G), \chi(G[W]) \leq h(n)\}$$

be the maximum number of vertices we can colour with h colours. Consider the vertex exposure martingale: modifying the edges from a single vertex can change $s(G)$ by at most 1, so the Azuma-Hoeffding inequality implies that, for $t > 0$,

$$\mathbb{P}(|s(G) - \mathbb{E}s(G)| > t) \leq 2e^{-t^2/n}.$$

In particular, for n sufficiently large,

$$\mathbb{P}(|s(G) - \mathbb{E}s(G)| > \sqrt{\omega(n)n}) < \frac{1}{\omega(n)}. \quad (3)$$

It follows from (2) that $\mathbb{P}(s(G) = n) > 1/\omega(n)$. Thus (3) implies that $\mathbb{E}(s(G)) \geq n - \sqrt{\omega(n)n}$ and therefore $\mathbb{P}(s(G) < n - 2\sqrt{\omega(n)n}) < 1/\omega(n)$. Let $W \subset V(G)$ be a subset of maximal size such that $\chi(G[W]) \leq h(n)$, and let $U = V(G) \setminus W$. We claim that, with probability $1 - o(1)$, $\chi(G[U]) \leq \omega(n)\sqrt{n}/\log n$. Since $\chi(G) \leq \chi(G[W]) + \chi(G[U])$, (1) follows immediately.

The claim follows simply from Lemma 2. Indeed, $|U| \leq 2\sqrt{\omega(n)n}$ with probability $1 - o(1)$. Let us greedily remove independent sets of maximal size from U , giving a new colour to each, until $n^{1/3}$ vertices remain (and give new colours to each of these). The lemma (applied to $\overline{G}$) implies that, with probability $1 - o(1)$, we use at most $O(2\sqrt{\omega(n)n}/c(p) \log n + n^{1/3})$ colours, which is bounded by $\omega(n)\sqrt{n}/\log n$ for sufficiently large n . $\square$

References

- [1] D. Achlioptas and A. Naor, The two possible values of the chromatic number of a random graph, *Ann. Math.* **162** (2005), 1335–1351
- [2] N. Alon and M. Krivelevich, The concentration of the chromatic number of random graphs, *Combinatorica* **17** (1997), 303–313
- [3] B. Bollobás, *Random Graphs*, Cambridge Studies in Advanced Mathematics **73**, second edition, Cambridge Univ. Press, Cambridge, 2001
- [4] B. Bollobás, The chromatic number of random graphs, *Combinatorica* **8** (1988), 49–55
- [5] B. Bollobás, How sharp is the concentration of the chromatic number?, *Combinatorics, Probability and Computing* **13** (2004) 115–117
- [6] A. Coja-Oghlan, K. Panagiotou, A. Steger, On the chromatic number of random graphs, *J. Comb. Theory, Ser. B*, to appear
- [7] T. Łuczak, A note on the sharp concentration of the chromatic number of random graphs, *Combinatorica* **11** (1991), 295–297

- [8] T. Łuczak, The chromatic number of random graphs, *Combinatorica* **11** (1991), 45–54
- [9] C. McDiarmid, On the chromatic number of random graphs, *Random Structures and Algorithms* **1** (1990), 435–442
- [10] E. Shamir and J. Spencer, Sharp concentration of the chromatic number on random graphs $G_{n,p}$, *Combinatorica* **7** (1987), 121–129